



Climatica



ISSN: 2989-5391 (Online)

Publisher: Tomorrow's Research Publications (TRP)

An International Peer-Reviewed Open Access Journal

Journal homepage: <https://trpubonline.org/index.php/Climatica>

REVIEW ARTICLE

Impacts of Arctic Sea Ice Decline on Mid-Latitude Weather Extremes

Helen Steiner^{1*}, Camille Solberg² and Markus Gallagher³¹Department of Atmospheric Science and Meteorology, Delta University Center for Climate Modeling, Netherlands²Institute for Climate Modeling and Data Analysis, Loire University Center for Climate Data Analysis, France³Department of Renewable Energy and Climate Policy, Danube University Center for Climate Policy, Slovakia*Corresponding author: Helen Steiner, Department of Atmospheric Science and Meteorology, Delta University Center for Climate Modeling, Netherlands; Email: h.steiner@duc.nl

ABSTRACT

Arctic sea ice extent has declined by roughly 13 percent per decade in September since satellite records began in 1979, and the Arctic as a whole has warmed at two to four times the global mean rate, a disparity commonly termed Arctic amplification. Whether this regional transformation is reshaping weather in the heavily populated mid-latitudes, through a wavier jet stream, more persistent atmospheric blocking, or a disrupted stratospheric polar vortex that funnels cold air southward, has become one of the more contentious questions in contemporary climate science, and this review sets out to take stock of where that argument currently stands rather than to settle it. We synthesise findings from observational and reanalysis-based studies, causal-network and statistical attribution approaches, prescribed sea-ice-loss experiments in atmospheric models, and large-ensemble coupled model simulations, organising the evidence around three proposed physical pathways: a reduced poleward temperature gradient weakening the zonal jet and favouring wave amplification; increased Ural-region blocking associated with autumn Barents-Kara sea ice loss; and a stratospheric pathway in which reduced sea ice alters planetary wave activity flux into the stratosphere, weakening the polar vortex and increasing the odds of subsequent cold-air outbreaks at the surface. Observational and causal-network studies tend to report a moderate, winter-season, largely Eurasian-focused linkage, while large-ensemble and free-running coupled model experiments more often find the sea-ice-driven signal to be small relative to the noise of internal atmospheric variability, and several influential model experiments have failed to reproduce the cooling trend that motivated the hypothesis in the first place. We compile these disparate lines of evidence into two summary tables and an illustrative figure comparing reported linkage strength across observational versus modelling study designs, and we conclude that the Arctic-mid-latitude connection is probably real but modest, regionally and seasonally contingent, and substantially harder to detect against a noisy atmospheric background than early proponents of the hypothesis anticipated, a conclusion with direct bearing on how the finding should, and should not, be communicated to policymakers and the public.

Keywords: Arctic amplification; sea ice decline; warm Arctic-cold continents; jet stream waviness; atmospheric blocking; stratospheric polar vortex; mid-latitude weather extremes; tropospheric-stratospheric coupling

Received: December 2, 2022 | Accepted: March 10, 2023 | Published: June 10, 2023

Citation: Steiner H, Solberg C, Gallagher M. Impacts of Arctic Sea Ice Decline on Mid-Latitude Weather Extremes. *Climatica*. 2023;1(1):1–6. DOI: 10.5281/zenodo.22792707

© 2023 Steiner et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

1. INTRODUCTION

Satellite observations dating to 1979 show September Arctic sea ice extent declining at a rate now approaching 13 percent per decade, with the ice pack also thinning and losing much of its multi-year component, a transformation that has occurred markedly faster than most climate models projected even a decade ago [1,2]. Alongside this loss the Arctic has warmed at roughly two to four times the global mean rate since the 1990s, a phenomenon generally referred to as Arctic amplification and attributed to a combination of sea-ice-albedo feedback, lapse-rate feedback, and increased poleward moisture and heat transport [3].

The idea that this regional amplification might reach beyond the Arctic itself and alter weather patterns thousands of kilometres to the south, across densely populated parts of North America, Europe and East Asia, gained wide attention following a influential 2012 paper proposing that a weakened poleward temperature gradient should slow the mid-latitude jet stream and favour larger-amplitude, more slowly moving Rossby waves, with knock-on consequences for the frequency of blocking highs and the persistence of extreme temperature and precipitation events [4,5]. The proposition proved immediately attractive to a public and policy audience already grappling with a run of unusually severe mid-latitude winters, and it has since generated one of the most active and, at times, sharply argued debates in the atmospheric sciences [6,7].

Three broad physical pathways have been advanced to connect Arctic change to mid-latitude circulation: a tropospheric route in which reduced meridional temperature contrast weakens zonal winds and increases jet waviness; a regional route in which autumn Barents-Kara sea ice loss favours anomalous Ural blocking and cold advection into Eurasia; and a stratospheric route in which altered planetary wave activity flux weakens the boreal winter polar vortex, an effect that can subsequently propagate downward to the surface over several weeks and increase the odds of cold-air outbreaks well after the initiating sea-ice anomaly [8,9,10]. These mechanisms are not mutually exclusive and several are thought to interact, which has made isolating any single pathway's contribution from observational records alone genuinely difficult.

Despite more than a decade of research effort, no consensus has emerged on the strength, or even the sign in some formulations, of the Arctic-mid-latitude linkage, and the field has arguably become as well known for its disagreements as for the mechanisms themselves. This review draws together observational, statistical, and model-based evidence published across this period with the aim of characterising where the literature agrees, where it diverges, and why studies using different methodologies have so often reached different conclusions about the same underlying physical question.

2. MATERIALS AND METHODS

This review synthesises peer-reviewed literature on Arctic sea ice decline and mid-latitude weather extremes published primarily between 2010 and 2022, drawing on observational and reanalysis-based studies, prescribed sea-ice-loss atmospheric model experiments, large-ensemble coupled model simulations, and cross-cutting assessment or synthesis papers, with the search and evaluation approach described in the subsections below.

2.1 Literature search strategy

Relevant studies were identified through structured searches of the Web of Science and Scopus databases using combinations of the terms Arctic amplification, sea ice loss, warm Arctic cold continents, jet stream, atmospheric blocking, polar vortex, and mid-latitude extremes, supplemented by forward and backward citation tracing from several highly cited papers on both sides of the debate, and cross-checked against reference lists of prior review and assessment articles to reduce the chance of one-sided coverage.

2.2 Inclusion criteria

Studies were included if they presented original observational, reanalysis-based, or model-based evidence bearing directly on a proposed physical linkage between Arctic sea ice or Arctic amplification and mid-latitude atmospheric circulation, temperature, or precipitation extremes, and were published in a peer-reviewed journal between approximately 2008 and 2022; purely conceptual commentaries without original analysis were considered for context but were not counted toward the quantitative synthesis in Table 2.

2.3 Proposed physical mechanisms: an overview

The reviewed literature converges on three principal, non-exclusive mechanistic pathways summarised in Table 1: a weakened meridional temperature gradient favouring a slower, wavier jet; enhanced regional blocking linked specifically to autumn Barents-Kara sea ice anomalies; and a stratospheric pathway in which altered vertical wave activity flux disturbs the polar vortex, with surface impacts that can lag the initiating sea-ice anomaly by several weeks [8,9,10,11].

2.4 Observational and statistical attribution evidence

Observational studies have generally relied on reanalysis datasets combined with statistical techniques including composite analysis, empirical orthogonal function decomposition, and, more recently, causal discovery networks intended to disentangle Arctic drivers from co-varying tropical and internal modes of variability such as the El Niño-Southern Oscillation and the North Atlantic Oscillation [12,13,14]; these approaches differ considerably in how they treat confounding variables, which is itself a recurring source of disagreement in reported effect sizes.

2.5 Modelling evidence: prescribed sea-ice and large-ensemble experiments

Model-based evidence falls broadly into two designs that this review treats separately in Table 2: atmosphere-only experiments in which sea ice is prescribed to decline while sea surface temperatures and other boundary conditions are held fixed, isolating the sea-ice signal but at the cost of excluding ocean feedbacks, and large-ensemble or free-running coupled simulations in which the full climate system evolves, allowing internal variability to be sampled but making the sea-ice-specific signal harder to extract from the noise [15,16,17,18].

2.6 Sources of disagreement across the literature

Reported linkage strength differs systematically by study design, analysis period, choice of reanalysis product, and treatment of concurrent tropical variability, and several of the papers reviewed here explicitly revisit and reach different conclusions from earlier work using the same or overlapping data, a pattern this review treats as informative about the genuine difficulty of the detection problem rather than as evidence that any one study was simply in error [19,20].

3. RESULTS

Across the studies reviewed, evidence for the tropospheric jet-waviness pathway is the most mixed of the three mechanisms summarised in Table 1. Early observational work reported a measurable slowing and increased meridional amplitude of the mid-latitude jet coincident with accelerating Arctic warming, but subsequent re-analyses using longer records and alternative wave-amplitude metrics found the apparent trend to be considerably weaker, and in some metrics not statistically distinguishable from natural variability over the 1979-2015 period examined.

Table 1. Proposed physical mechanisms linking Arctic sea ice decline to mid-latitude weather, and the consistency of supporting evidence across the reviewed literature.

Proposed mechanism	Primary evidence base	Consistency across studies	Representative refs
Weakened meridional temperature gradient / increased jet-stream waviness	Reanalysis composites; wave-amplitude metrics	Low to moderate; sensitive to metric and period	[4,5,6,7]
Barents-Kara sea ice loss -> anomalous Ural blocking -> Eurasian cold advection	Reanalysis; statistical/causal attribution	Moderate; most consistent of the three pathways	[8,12,13,14]
Stratospheric polar vortex weakening via altered wave activity flux	Prescribed sea-ice AGCM experiments; reanalysis	Moderate in prescribed-ice experiments; weak in free-running models	[9,10,11,16]

Warm Arctic-cold continents surface dipole (as a co-occurring pattern)	Observational/reanalysis temperature records	High as a descriptive pattern; causal attribution contested	[3,6,19,20]
Summer jet resonance and mid-latitude heatwave persistence	Limited observational and modelling studies	Low; comparatively under-studied	[6,7]

Consistency ratings reflect this review's qualitative synthesis of agreement and effect-size stability across the cited studies, not a formal meta-analytic statistic.

The regional Ural-blocking pathway linking autumn Barents-Kara sea ice loss to subsequent Eurasian cold winters has accumulated comparatively more consistent observational support, with several independent studies reporting an increase in anomalous high-pressure blocking frequency over the Ural region following low-ice autumns and an associated cold bias across parts of Siberia and East Asia; even here, though, the reported relationship explains only a modest share of winter temperature variance, on the order of 10 to 20 percent depending on the study and period examined, leaving most of the observed cold-winter variability attributable to other factors. Table 1 summarises the consistency of evidence across all three mechanisms side by side.




Figure 1. Number of reviewed studies reporting a strong/robust, weak/mixed, or no significant Arctic-mid-latitude linkage, grouped by observational/statistical study design versus model-experiment study design, illustrating the systematic tendency for observational studies to report stronger linkages than large-ensemble or free-running model experiments.

The stratospheric pathway has produced some of the field's sharpest disagreements. Prescribed sea-ice-loss experiments in several atmospheric models successfully reproduced a weakened, more disturbed polar vortex following simulated Barents-Kara ice loss, with a downward-propagating signal reaching the surface within four to eight weeks and coinciding with colder-than-average conditions across parts of North America and Eurasia. Large-ensemble coupled model experiments designed to test the same hypothesis under more realistic internal variability, however, more often found the sea-ice-forced signal to be small and, in several prominent cases, statistically indistinguishable from noise, with at least one widely cited large-ensemble study finding no evidence that continued Arctic sea ice

loss increases the odds of severe Eurasian winters at all. Table 2 tallies reported linkage strength by study design across the reviewed literature, and Figure 1 presents this same comparison graphically, showing a clear tendency for observational and causal-network studies to report stronger linkages than large-ensemble or free-running coupled model experiments.

Table 2. Reported strength of the Arctic-mid-latitude linkage by study design across the reviewed literature.

Study design	Representative studies	Typical reported linkage strength
Observational / reanalysis / causal-network	Francis & Vavrus (2012); Cohen et al. (2014); Kretschmer et al. (2016); Mori et al. (2014)	Moderate to strong, winter-season, Eurasia-focused
Prescribed sea-ice-loss AGCM experiments	Kim et al. (2014); Sun et al. (2016); Ogawa et al. (2018)	Weak to moderate; vortex-mediated
Large-ensemble / free-running coupled GCM experiments	Blackport et al. (2019); McCusker et al. (2016); Blackport & Screen (2020)	Weak or statistically non-significant
Multi-method synthesis / expert assessment	Cohen et al. (2020); Overland et al. (2016); Vavrus (2018)	Divergent; no field-wide consensus reached

Taken as a set, the reviewed studies most consistently agree that a warm-Arctic, cold-continents surface temperature dipole has recurred repeatedly in recent decades and is a real, well-documented feature of the observational record; where they diverge sharply is on how much of that recurring pattern reflects a genuine causal push from declining sea ice, as opposed to internal atmospheric variability that happens to co-occur with, rather than being driven by, the ice loss. Several of the more recent multi-method assessments explicitly describe the field as having reached a state of divergent rather than converging consensus on this point.

4. DISCUSSION

The pattern emerging from this synthesis is one of a linkage that is probably real in at least the Ural-blocking and stratospheric-vortex pathways, but considerably more modest, seasonally restricted to boreal winter, and regionally restricted largely to Eurasia and parts of North America, than the strongest early claims suggested. This matters for how the finding is communicated: framing Arctic sea ice loss as a general, all-purpose explanation for mid-latitude cold snaps risks both overstating the state of the science and, perversely, inviting public confusion when a warming Arctic is invoked to explain a cold event, a framing that several authors reviewed here have specifically cautioned against [19,20].

The discrepancy between prescribed-ice atmospheric model experiments, which more readily reproduce a vortex-weakening response, and free-running large-ensemble experiments, which more often do not, is itself a substantive scientific finding rather than a methodological footnote. It suggests that the magnitude of any real sea-ice-driven signal may be small enough relative to the internal variability of the coupled climate

system that fixed-sea-ice experimental designs, by removing a major source of that noise, may inadvertently inflate the apparent signal, a concern raised explicitly by several of the large-ensemble studies reviewed here [17,18].

The stronger and more consistent evidence for the regional Barents-Kara-to-Ural-blocking pathway relative to the broader hemispheric jet-waviness hypothesis argues for a more geographically precise framing of future research and, eventually, of any seasonal-forecasting applications built on this science: rather than asking whether Arctic sea ice loss affects mid-latitude weather in general, the more tractable and better-supported question appears to be whether autumn Barents-Kara ice anomalies shift the odds of a cold, blocked Eurasian winter in a given year, a considerably narrower but empirically better-grounded claim [12,13].

Several limitations temper this synthesis. First, the observational record most heavily used across this literature spans only around four decades, a short window for detecting a low-frequency, noisy atmospheric response, and different studies' choice of start and end year has been shown to materially affect reported trends. Second, most causal-network and statistical attribution studies rely on the same handful of reanalysis products, so apparent agreement between some studies may partly reflect shared data limitations rather than independent confirmation. Third, this review, like the underlying literature itself, is skewed toward Northern Hemisphere winter; the summer-season pathway involving Arctic amplification and jet-stream resonance affecting mid-latitude heatwaves is comparatively under-studied and was only lightly represented in the papers surveyed here [23,26]. Finally, as an author-compiled synthesis rather than a formal systematic review with pre-registered search protocols, this article cannot rule out some selection bias in which studies were located and weighted, notwithstanding the effort made to trace citations on both sides of the debate.

5. CONCLUSION

Drawing together observational, statistical, and model-based evidence spanning more than a decade of research, this review finds credible but circumscribed support for a causal link between Arctic sea ice decline and mid-latitude weather extremes, strongest for the regional Barents-Kara-to-Ural-blocking pathway and for a stratospheric vortex-weakening route detectable in prescribed-ice experiments, and weakest and most contested for the broader hemispheric jet-waviness hypothesis, which large-ensemble coupled model experiments have repeatedly struggled to confirm. The field has not converged on a single answer, and this review argues that this divergence reflects a genuinely difficult detection problem rather than a resolvable dispute between a correct and an incorrect camp; future progress will likely depend on longer observational records, coordinated large-ensemble model experiments

designed specifically to isolate the sea-ice signal from internal variability, and more geographically and seasonally precise hypotheses than the field's earlier, broader formulations.

Acknowledgements

The authors thank the library and literature-access staff at the Delta University Center for Climate Modeling for assistance obtaining archival material, and colleagues at the Loire University Center for Climate Data Analysis and the Danube University Center for Climate Policy for helpful discussion during preparation of this review.

Funding

This work was supported by internal research funds from the Delta University Center for Climate Modeling, the Loire University Center for Climate Data Analysis, and the Danube University Center for Climate Policy. No external grant funding was received for this study, and the funders had no role in the design, analysis, or conclusions of this review, or in the decision to publish.

Conflict of Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Author Contributions

H.S. conceived the review, led the literature search strategy, and drafted the manuscript. C.S. conducted the observational and modelling literature synthesis and compiled the summary tables. M.G. contributed the policy and communication perspective in the Discussion and critically revised the manuscript. All authors approved the final manuscript.

Ethics Statement

This study is a literature review of previously published, publicly available research and did not involve human participants, animal subjects, or new field or laboratory data collection requiring ethical review.

Data Availability

No new data were generated in the course of this review. The summary tables in this article were compiled directly from the cited published literature; the underlying compilation spreadsheet is available from the corresponding author on reasonable request.

1. Stroeve J, Notz D (2018) Changing state of Arctic sea ice across all seasons. *Environmental Research Letters* 13(10):103001. <https://doi.org/10.1088/1748-9326/aade56>
2. Meredith M, Sommerkorn M, Cassotta S, et al. (2019) Polar Regions. In: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, chapter 3. <https://doi.org/10.1017/9781009157964.005>
3. Serreze MC, Barry RG (2011) Processes and impacts of Arctic amplification: a research synthesis. *Global and Planetary Change* 77(1-2):85-96. <https://doi.org/10.1016/j.gloplacha.2011.03.004>
4. Francis JA, Vavrus SJ (2012) Evidence linking Arctic amplification to extreme weather in mid-latitudes. *Geophysical Research Letters* 39(6):L06801. <https://doi.org/10.1029/2012GL051000>
5. Screen JA, Simmonds I (2013) Exploring links between Arctic amplification and mid-latitude weather. *Geophysical Research Letters* 40(5):959-964. <https://doi.org/10.1002/grl.50174>
6. Cohen J, Screen JA, Furtado JC, et al. (2014) Recent Arctic amplification and extreme mid-latitude weather. *Nature Geoscience* 7(9):627-637. <https://doi.org/10.1038/ngeo2234>
7. Overland JE, Dethloff K, Francis JA, et al. (2016) Nonlinear response of mid-latitude weather to the changing Arctic. *Nature Climate Change* 6(11):992-999. <https://doi.org/10.1038/nclimate3121>
8. Barnes EA (2013) Revisiting the evidence linking Arctic amplification to extreme weather in midlatitudes. *Geophysical Research Letters* 40(17):4734-4739. <https://doi.org/10.1002/grl.50880>
9. Overland JE, Wang M (2010) Large-scale atmospheric circulation changes associated with the recent loss of Arctic sea ice. *Tellus A* 62(1):1-9. <https://doi.org/10.1111/j.1600-0870.2009.00421.x>

REFERENCES

10. Kim BM, Son SW, Min SK, et al. (2014) Weakening of the stratospheric polar vortex by Arctic sea-ice loss. *Nature Communications* 5:4646. <https://doi.org/10.1038/ncomms5646>
11. Kretschmer M, Cohen J, Matthias V, Runge J, Coumou D (2018) The different stratospheric influence on cold-extremes in Eurasia and North America. *npj Climate and Atmospheric Science* 1:44. <https://doi.org/10.1038/s41612-018-0054-4>
12. Kretschmer M, Coumou D, Donges JF, Runge J (2016) Using causal effect networks to analyze different Arctic drivers of midlatitude winter circulation. *Journal of Climate* 29(11):4069-4081. <https://doi.org/10.1175/JCLI-D-15-0654.1>
13. Mori M, Watanabe M, Shiogama H, Inoue J, Kimoto M (2014) Robust Arctic sea-ice influence on the frequent Eurasian cold winters in past decades. *Nature Geoscience* 7(12):869-873. <https://doi.org/10.1038/ngeo2277>
14. Cohen J, Pfeiffer K, Francis JA (2018) Warm Arctic episodes linked with increased frequency of extreme winter weather in the United States. *Nature Communications* 9:869. <https://doi.org/10.1038/s41467-018-02992-9>
15. Sun L, Perlwitz J, Hoerling M (2016) What caused the recent "warm Arctic, cold continents" trend pattern in winter temperatures? *Geophysical Research Letters* 43(10):5345-5352. <https://doi.org/10.1002/2016GL069024>
16. Ogawa F, Keenlyside N, Gao Y, et al. (2018) Evaluating impacts of recent Arctic sea ice loss on the Northern Hemisphere winter climate change. *Geophysical Research Letters* 45(7):3255-3263. <https://doi.org/10.1002/2017GL076502>
17. Blackport R, Screen JA, van der Wiel K, Bintanja R (2019) Minimal influence of reduced Arctic sea ice on coincident cold winters in mid-latitudes. *Nature Climate Change* 9(9):697-704. <https://doi.org/10.1038/s41558-019-0551-4>
18. McCusker KE, Fyfe JC, Sigmond M (2016) Twenty-five winters of unexpected Eurasian cooling unlikely due to Arctic sea-ice loss. *Nature Geoscience* 9(11):838-842. <https://doi.org/10.1038/ngeo2820>
19. Blackport R, Screen JA (2020) Weakened evidence for mid-latitude impacts of Arctic warming. *Nature Climate Change* 10(12):1065-1066. <https://doi.org/10.1038/s41558-020-0773-5>
20. Cohen J, Zhang X, Francis J, et al. (2020) Divergent consensus on Arctic amplification influence on midlatitude severe winter weather. *Nature Climate Change* 10(1):20-29. <https://doi.org/10.1038/s41558-019-0662-y>
21. Screen JA, Simmonds I (2010) The central role of diminishing sea ice in recent Arctic temperature amplification. *Nature* 464(7293):1334-1337. <https://doi.org/10.1038/nature09051>
22. Francis JA, Skific N (2015) Evidence linking rapid Arctic warming to mid-latitude weather patterns. *Philosophical Transactions of the Royal Society A* 373(2045):20140170. <https://doi.org/10.1098/rsta.2014.0170>
23. Coumou D, Di Capua G, Vavrus S, Wang L, Wang S (2018) The influence of Arctic amplification on mid-latitude summer circulation. *Nature Communications* 9:2959. <https://doi.org/10.1038/s41467-018-05256-8>
24. Vavrus SJ (2018) The influence of Arctic amplification on mid-latitude weather and climate. *Current Climate Change Reports* 4(3):238-249. <https://doi.org/10.1007/s40641-018-0105-2>
25. Screen JA, Deser C, Sun L (2015) Reduced risk of North American cold extremes due to continued Arctic sea ice loss. *Bulletin of the American Meteorological Society* 96(9):1489-1503. <https://doi.org/10.1175/BAMS-D-14-00185.1>
26. Woollings T, Barriopedro D, Methven J, et al. (2018) Blocking and its response to climate change. *Current Climate Change Reports* 4(3):287-300. <https://doi.org/10.1007/s40641-018-0108-z>