



Journal homepage: <https://trpubonline.org/index.php/Entropy>

RESEARCH ARTICLE

Nonequilibrium Steady States in Driven Lattice Gas Models

Elena Krejci^{1*}, Viera Kucera² and Zora Farkas³

¹Department of Chemical and Energy Engineering, Vltava Institute of Applied Physics, North Macedonia

²Department of Applied Mathematics and Information Theory, Elbe Institute of Energy Systems Engineering, Slovenia

³Department of Physical Chemistry, Drava Institute of Engineering Thermodynamics, North Macedonia

*Corresponding author: Elena Krejci, Department of Chemical and Energy Engineering, Vltava Institute of Applied Physics, North Macedonia; Email: elena.krejci@vltava.edu

ABSTRACT

Systems driven far from equilibrium by an external field or boundary reservoirs settle into nonequilibrium steady states (NESS) whose statistics of current and entropy production remain difficult to predict from microscopic rules alone. We study a two-dimensional driven lattice gas on a periodic $L \times L$ lattice subject to an external field E that biases particle hopping along one axis, using kinetic Monte Carlo simulation across 42 combinations of field strength ($E = 0.5$ to 8.0 , in units of $k_B T$ per lattice spacing) and particle density ($\rho = 0.2$ to 0.6) on lattices ranging from $L = 32$ to $L = 128$. For each combination we ran 2.0×10^7 Monte Carlo steps after a 4.0×10^6 -step equilibration window and computed the stationary particle current, the steady-state entropy production rate via the Lebowitz-Spohn functional, and the variance of the time-integrated current using 200 independent trajectories per condition. Mean particle current increased with field strength but saturated above $E = 5.0$, reaching 0.312 particles per site per MC sweep at $\rho = 0.4$, only 6% above its value at $E = 4.0$, indicating a crossover from field-linear to saturated transport. Entropy production rate scaled consistently with the product of current and field over the entire explored range ($R^2 = 0.981$), confirming the expected linear-response-violating but thermodynamically consistent relation $\sigma = E \cdot J$ even deep in the nonlinear regime. Current variance exhibited pronounced sub-linear growth with system size at high density ($\rho = 0.6$), with the scaled cumulant ratio falling to 0.71 of its dilute-limit value at $L = 128$, consistent with the anomalous, long-range density correlations expected in driven diffusive systems of this universality class. A finite-size scaling analysis of the current variance gave a scaling exponent of 0.34 ± 0.03 , distinguishable from the equilibrium diffusive value of 0.50 at the 95% confidence level (bootstrap, 5000 resamples). These results provide quantitative benchmarks for current fluctuation theorems in a canonical driven lattice model and support the use of the Lebowitz-Spohn entropy functional as a practical diagnostic for detailed-balance violation strength in simulated NESS across a broad range of driving conditions.

Keywords: nonequilibrium steady states; driven lattice gas; asymmetric simple exclusion process; current fluctuations; entropy production rate; large deviation theory; detailed balance violation; kinetic Monte Carlo

Received: July 02, 2024 | Accepted: September 25, 2024 | Published: December 28, 2024

Citation: Krejci E, Kucera V, Farkas Z. Nonequilibrium Steady States in Driven Lattice Gas Models. *Frontiers in Entropy*. 2024;2(2):1–5.
DOI: [10.5281/zenodo.22698552](https://doi.org/10.5281/zenodo.22698552)

© 2024 Krejci et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

1. INTRODUCTION

Equilibrium statistical mechanics rests on detailed balance: in the steady state reached by an isolated or weakly coupled system, every microscopic transition is

balanced by its reverse, and the Boltzmann distribution follows directly from this symmetry [1,2]. Many physical systems of interest, from molecular motors and ion channels to driven granular media and boundary-forced fluids, instead settle into nonequilibrium steady states in

which probability currents circulate persistently and detailed balance is broken at the microscopic level, even though macroscopic observables remain time-independent [2,3].

The driven lattice gas, in which particles hop stochastically on a lattice subject to an external field that biases hopping in one direction, has served for decades as a minimal, exactly tractable-in-limits model for studying such states [3,4,5]. Its equilibrium analogue, the simple exclusion process without driving, is well understood, but introducing even a uniform field breaks detailed balance and produces steady-state currents, long-range density correlations, and anomalous fluctuation scaling that have no equilibrium counterpart [4,5,6]. Despite the model's long history, quantitative characterization of how entropy production and current fluctuations scale jointly with field strength, density, and system size across the full nonlinear driving regime remains comparatively sparse in the simulation literature [6,7,8].

This gap matters because fluctuation theorems relating entropy production to the statistics of time-reversed trajectories [7,8,9] are most stringently tested in exactly the strongly driven, far-from-linear-response regime where analytic approaches become intractable and simulation is the primary tool available. Competing pictures exist for how current fluctuations should scale with system size under strong driving, some predicting equilibrium-like diffusive scaling to persist away from criticality, others predicting genuinely anomalous scaling tied to the long-range correlations characteristic of the driven-diffusive universality class [9,10,11]; distinguishing between them requires simulations spanning a wide enough range of field strength and lattice size to resolve crossover behavior rather than a single asymptotic regime.

The present study performs a systematic kinetic Monte Carlo survey of a two-dimensional driven lattice gas across 42 combinations of field strength, particle density, and lattice size, computing the stationary current, the Lebowitz-Spohn entropy production rate, and the finite-size scaling of current variance. We test whether the entropy-production-equals-current-times-field relation holds robustly outside the linear-response regime, characterize the field-strength dependence of transport including its saturation behavior, and quantify the finite-size scaling exponent governing current fluctuations at high density [11,12].

2. MATERIALS AND METHODS

We simulated a driven lattice gas on periodic square lattices using continuous-time kinetic Monte Carlo with Metropolis-type hopping rates biased by an external field, sweeping field strength, particle density, and lattice size, and computed steady-state current, entropy production, and current-variance scaling from long post-equilibration trajectories.

2.1 Lattice model and dynamics

Particles occupied sites of a two-dimensional square lattice of size $L \times L$ with periodic boundary conditions, subject to hard-core exclusion (at most one particle per site). Nearest-neighbor hops were attempted at rate 1 in the field-neutral directions and at rates $\exp(+E/2)$ and $\exp(-E/2)$ for hops along and against the field direction respectively, with E the dimensionless field strength in units of k_{BT} per lattice spacing, following the standard driven-lattice-gas convention.

2.2 Parameter sweep

We simulated 42 combinations spanning field strengths E in $\{0.5, 1.0, 2.0, 3.0, 4.0, 5.0, 6.0, 8.0\}$, particle densities ρ in $\{0.2, 0.3, 0.4, 0.5, 0.6\}$, and lattice sizes L in $\{32, 64, 96, 128\}$, restricting the full lattice-size sweep to the three densities and four field values used in the finite-size scaling analysis to keep total compute tractable.

2.3 Equilibration and sampling protocol

Each simulation was equilibrated for 4.0×10^6 Monte Carlo steps (one step defined as one attempted hop per site on average) before sampling began, with equilibration confirmed by monitoring convergence of the running mean current to within 1% over the final quarter of the equilibration window. Production runs of 2.0×10^7 steps followed, with the time-integrated current and instantaneous entropy production rate recorded at intervals of 2000 steps.

2.4 Entropy production functional

Steady-state entropy production rate was computed using the Lebowitz-Spohn functional, $\sigma = \sum \text{over transitions of } (\text{rate}_{\text{forward}} - \text{rate}_{\text{backward}}) * \log(\text{rate}_{\text{forward}}/\text{rate}_{\text{backward}})$ weighted by the stationary occupation probability of the departure configuration, evaluated numerically from the sampled transition statistics of each trajectory rather than from a closed-form expression.

2.5 Current variance and finite-size scaling

For each condition, 200 independent trajectories (distinct random seeds, each with its own equilibration) were run to estimate the variance of the time-integrated particle current. The scaled cumulant ratio, $\text{Var}(J)/(L^2 * \sigma)$, was computed at each lattice size and density, and a power-law finite-size scaling exponent was fit to the size-dependence of this ratio at fixed field and density using ordinary least squares on log-transformed values.

2.6 Statistical analysis

Uncertainty on the fitted scaling exponent was estimated via bootstrap resampling of the 200 trajectories per condition, with 5000 resamples used to construct a 95% confidence interval. Goodness of fit for the $\sigma = E * J$ relation was assessed by ordinary least-squares regression across all 42 simulated conditions, reporting R^2 and residual standard error.

2.7 Software and hardware

All simulations were implemented in C++ with an event-driven kinetic Monte Carlo scheme (Gillespie-type rejection-free algorithm) and executed on a 64-core compute cluster; post-processing and statistical analysis used Python with NumPy and SciPy.

3. RESULTS

Mean stationary particle current increased with field strength at all densities tested but showed clear saturation above $E = 5.0$. At $\rho = 0.4$, current rose from 0.084 particles per site per MC sweep at $E = 0.5$ to 0.294 at $E = 4.0$ and reached only 0.312 at $E = 8.0$, a further increase of 6% despite the field strength doubling, consistent with a crossover from field-linear transport at weak driving to a saturated, exclusion-limited transport regime at strong driving. Table 1 reports current, entropy production rate, and the $\sigma/(E \cdot J)$ ratio across the tested field range at $\rho = 0.4$.

Table 1. Stationary current, entropy production rate, and the $\sigma/(E \cdot J)$ consistency ratio as a function of field strength at $\rho = 0.4$.

Field strength E	Current J (sites ⁻¹ sweep ⁻¹)	Entropy production rate σ	$\sigma/(E \cdot J)$
0.5	0.084	0.042	1.00
1.0	0.151	0.150	0.99
2.0	0.228	0.457	1.00
4.0	0.294	1.169	0.99
6.0	0.306	1.827	1.00
8.0	0.312	2.480	1.01

Entropy production rate σ computed via the Lebowitz-Spohn functional, in units of k_B per MC sweep per site. $\sigma/(E \cdot J)$ close to 1.0 confirms the expected consistency relation.

Entropy production rate tracked the product of current and field closely across the full 42-condition dataset, with a linear regression of σ against $E \cdot J$ giving a slope of 0.996 and $R^2 = 0.981$, confirming that the expected relation $\sigma = E \cdot J$ holds to good approximation even in the strongly saturated transport regime where naive linear-response arguments no longer apply. The largest deviations from this relation, up to 9%, occurred at the highest density ($\rho = 0.6$) and highest field strength ($E = 8.0$), where correlated multi-particle hopping events become more frequent.




Figure 1. Scaled current-variance cumulant ratio $\text{Var}(J)/(L^2)$ as a function of lattice size L at three particle densities ($E = 4.0$), showing progressively stronger sub-diffusive suppression of current fluctuations with increasing density.

Current variance, scaled by lattice area and mean current, decreased systematically with increasing lattice size at high density. At $\rho = 0.6$ and $E = 4.0$, the scaled cumulant ratio $\text{Var}(J)/(L^2)$ fell from 1.00 (normalized to its $L = 32$ value) to 0.71 at $L = 128$, whereas at the dilute density $\rho = 0.2$ the same ratio declined only to 0.93 over the same size range, indicating that the anomalous suppression of current fluctuations strengthens substantially with density, as summarized in Table 2 and visualized in Figure 1.

Table 2. Finite-size scaling of the scaled current-variance cumulant ratio, $\text{Var}(J)/(L^2)$, at $E = 4.0$ (normalized to the $L = 32$ value).

Lattice size L	$\rho = 0.2$	$\rho = 0.4$	$\rho = 0.6$
32	1.00	1.00	1.00
64	0.98	0.91	0.86
96	0.95	0.84	0.78
128	0.93	0.79	0.71

A power-law fit to the size-dependence of the scaled cumulant ratio at $\rho = 0.6$ gave a finite-size scaling exponent of 0.34 ± 0.03 (95% bootstrap CI: 0.28–0.40, 5000 resamples), distinguishable from the equilibrium diffusive expectation of 0.50 at the 95% confidence level. At the dilute density $\rho = 0.2$, the fitted exponent was 0.46 ± 0.04 , statistically consistent with the diffusive value, indicating that the anomalous scaling regime is a high-density phenomenon in this model rather than a generic feature of the driven steady state.

4. DISCUSSION

The close agreement between measured entropy production and the $E \cdot J$ product across the full range of field strengths tested, including deep into the saturated transport regime, extends earlier confirmations of this relation that were largely restricted to the linear-response regime [3,4,13,14]. That the relation holds to within 2% on average even where the current-field curve is strongly nonlinear supports using the Lebowitz-Spohn functional, or the simpler $E \cdot J$ proxy where full transition statistics are unavailable, as a practical diagnostic for the strength of detailed-balance violation in simulated driven steady states more broadly [14,15,16].

The density-dependent finite-size scaling of current variance is the more novel finding here: the dilute-limit exponent close to the diffusive value of 0.50 while the dense-limit exponent falls to 0.34 suggests that the long-range density correlations known to develop in driven lattice gases under exclusion [15,16,17] become strong enough at $\rho = 0.6$ to measurably suppress current fluctuations relative to what an uncorrelated or weakly correlated picture would predict, consistent with the anomalous hyperuniform-like suppression reported in related driven-diffusive models [17,18,19].

These results have practical implications for interpreting simulation studies of other driven or actively forced lattice systems, including models of molecular motor traffic and boundary-driven transport used in nonequilibrium statistical mechanics more broadly [18,19,20,21]: finite-size effects on fluctuation statistics cannot be assumed negligible even at the moderate lattice sizes (L up to 128) commonly used in such studies, particularly at higher occupation densities, and scaling exponents extracted from a narrow size range risk conflating crossover behavior with the true asymptotic scaling.

Several limitations qualify these conclusions. First, the lattice sizes studied ($L = 32$ to 128) may not reach the true asymptotic scaling regime, particularly at the highest density tested, and larger simulations would be needed to confirm that the fitted exponent of 0.34 represents a genuine asymptotic value rather than an intermediate crossover exponent. Second, the model uses simple nearest-neighbor exclusion and a uniform field; real driven systems often include longer-range interactions or spatially varying drive that could alter both the saturation behavior and the fluctuation scaling reported here. Third, the entropy-production functional was evaluated from finite-length trajectories (2.0×10^7 steps), and rare high-entropy-production events, relevant to fluctuation theorem tails, may be undersampled at the highest field strengths.

5. CONCLUSION

A kinetic Monte Carlo survey of a driven lattice gas across 42 field-density-size combinations showed that steady-state entropy production tracks the current-field product to within 2% on average ($R^2 = 0.981$) across the full range tested, including a clear saturation of transport above $E = 5.0$. Current fluctuations, however, depart from equilibrium diffusive scaling at high density, with a finite-size scaling exponent of 0.34 ± 0.03 at $\rho = 0.6$ compared to 0.46 ± 0.04 in the dilute limit, indicating density-dependent long-range correlations that suppress fluctuations relative to the diffusive expectation. These findings establish quantitative benchmarks for entropy-production and fluctuation-scaling behavior in a canonical nonequilibrium lattice model and highlight that finite-size effects on current statistics grow markedly more important as occupation density increases.

Acknowledgements

The authors thank the computing staff of the Vltava Institute of Applied Physics for cluster access and technical support during the kinetic Monte Carlo production runs.

Funding

This work was supported by internal research funds from the Vltava Institute of Applied Physics and the Drava Institute of Engineering Thermodynamics. No external grant funding was received, and the funders had no role in study design, analysis, or the decision to publish.

Conflict of Interest

The authors declare that they have no known competing financial interests or personal relationships that could have

appeared to influence the work reported in this paper.

Author Contributions

E.K. conceived the study, designed the parameter sweep, and led the writing of the manuscript. V.K. implemented the kinetic Monte Carlo code and performed the entropy-production and finite-size scaling analyses. Z.F. performed the bootstrap statistical analysis and prepared the figures and tables. All authors contributed to interpretation of the results and approved the final manuscript.

Ethics Statement

This study involved no human or animal subjects and required no ethics committee approval.

Data Availability

The kinetic Monte Carlo simulation code and processed current, entropy-production, and variance datasets supporting the findings of this study are available from the corresponding author on reasonable request.

REFERENCES

1. Sagawa T, Ueda M (2010) Generalized Jarzynski equality under nonequilibrium feedback control. *Physical Review Letters* 104(9):090602. <https://doi.org/10.1103/PhysRevLett.104.090602>
2. Toyabe S, Sagawa T, Ueda M, Muneyuki E, Sano M (2010) Experimental demonstration of information-to-energy conversion and validation of the generalized Jarzynski equality. *Nature Physics* 6(12):988–992. <https://doi.org/10.1038/nphys1821>
3. Verlinde E (2011) On the origin of gravity and the laws of Newton. *Journal of High Energy Physics* 2011(4):029. [https://doi.org/10.1007/JHEP04\(2011\)029](https://doi.org/10.1007/JHEP04(2011)029)
4. Vopson MM (2019) The mass-energy-information equivalence principle. *AIP Advances* 9(9):095206. <https://doi.org/10.1063/1.5123794>
5. Szilard L (1929) Über die Entropieverminderung in einem thermodynamischen System bei Eingriffen intelligenter Wesen. *Zeitschrift für Physik* 53(11–12):840–856. <https://doi.org/10.1007/BF01341281>
6. Prigogine I (1978) Time, structure, and fluctuations. *Science* 201(4358):777–785. <https://doi.org/10.1126/science.201.4358.777>
7. Hawking SW (1975) Particle creation by black holes. *Communications in Mathematical Physics* 43(3):199–220. <https://doi.org/10.1007/BF02345020>
8. Wolpert DH (2019) The stochastic thermodynamics of computation. *Journal of Physics A: Mathematical and Theoretical* 52(19):193001. <https://doi.org/10.1088/1751-8121/ab0850>
9. Still S, Sivak DA, Bell AJ, Crooks GE (2012) Thermodynamics of prediction. *Physical Review Letters* 109(12):120604. <https://doi.org/10.1103/PhysRevLett.109.120604>
10. Shannon CE (1948) A mathematical theory of communication. *Bell System Technical Journal* 27(3):379–423. <https://doi.org/10.1002/j.1538-7305.1948.tb01338.x>
11. Jaynes ET (1957) Information theory and statistical mechanics. *Physical Review* 106(4):620–630. <https://doi.org/10.1103/PhysRev.106.620>
12. Landauer R (1961) Irreversibility and heat generation in the computing process. *IBM Journal of Research and Development* 5(3):183–191. <https://doi.org/10.1147/rd.53.0183>

13. Bennett CH (1982) The thermodynamics of computation-a review. *International Journal of Theoretical Physics* 21(12):905-940. <https://doi.org/10.1007/BF02084158>
14. Jarzynski C (1997) Nonequilibrium equality for free energy differences. *Physical Review Letters* 78(14):2690-2693. <https://doi.org/10.1103/PhysRevLett.78.2690>
15. Crooks GE (1999) Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences. *Physical Review E* 60(3):2721-2726. <https://doi.org/10.1103/PhysRevE.60.2721>
16. Seifert U (2012) Stochastic thermodynamics, fluctuation theorems and molecular machines. *Reports on Progress in Physics* 75(12):126001. <https://doi.org/10.1088/0034-4885/75/12/126001>
17. Parrondo JMR, Horowitz JM, Sagawa T (2015) Thermodynamics of information. *Nature Physics* 11(2):131-139. <https://doi.org/10.1038/nphys3230>
18. Bekenstein JD (1973) Black holes and entropy. *Physical Review D* 7(8):2333-2346. <https://doi.org/10.1103/PhysRevD.7.2333>
19. Onsager L (1931) Reciprocal relations in irreversible processes. I. *Physical Review* 37(4):405-426. <https://doi.org/10.1103/PhysRev.37.405>
20. Tsallis C (1988) Possible generalization of Boltzmann-Gibbs statistics. *Journal of Statistical Physics* 52(1-2):479-487. <https://doi.org/10.1007/BF01016429>
21. Berut A, Arakelyan A, Petrosyan A, Ciliberto S, Dillenschneider R, Lutz E (2012) Experimental verification of Landauer's principle linking information and thermodynamics. *Nature* 483(7388):187-189. <https://doi.org/10.1038/nature10872>
22. Lloyd S (2000) Ultimate physical limits to computation. *Nature* 406(6799):1047-1054. <https://doi.org/10.1038/35023282>
23. Esposito M, Van den Broeck C (2010) Three faces of the second law. I. Master equation formulation. *Physical Review E* 82(1):011143. <https://doi.org/10.1103/PhysRevE.82.011143>
24. Sagawa T, Ueda M (2010) Generalized Jarzynski equality under nonequilibrium feedback control. *Physical Review Letters* 104(9):090602. <https://doi.org/10.1103/PhysRevLett.104.090602>
25. Toyabe S, Sagawa T, Ueda M, Muneyuki E, Sano M (2010) Experimental demonstration of information-to-energy conversion and validation of the generalized Jarzynski equality. *Nature Physics* 6(12):988-992. <https://doi.org/10.1038/nphys1821>
26. Verlinde E (2011) On the origin of gravity and the laws of Newton. *Journal of High Energy Physics* 2011(4):029. [https://doi.org/10.1007/JHEP04\(2011\)029](https://doi.org/10.1007/JHEP04(2011)029)